



Australian Government
Australian Fisheries Management Authority




Commonwealth Small Pelagic Fishery: Status Summary Report 2019

Report to the Australian Fisheries Management Authority

T. M. Ward and G. L. Grammer

RR2019/0803 February 2021



Contents

<i>About this document</i>	3
<i>Acknowledgements</i>	4
Overview	5
Jack Mackerel	5
Blue Mackerel.....	6
Redbait.....	6
Australian Sardine.....	7
Summary.....	7

About this document

This publication may be cited as:

Ward, T. M. and Grammer, G. L. (2021). Commonwealth Small Pelagic Fishery: Status Summary Report 2019. Report to the Australian Fisheries Management Authority. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2020/000209-1. SARDI Research Report Series No. 1057. 7pp.

South Australian Research and Development Institute

SARDI Aquatic Sciences

2 Hamra Avenue

West Beach SA 5024

Telephone: (08) 8207 5400

Facsimile: (08) 8207 5415

<http://www.pir.sa.gov.au/research>

DISCLAIMER

The authors warrant that they have taken all reasonable care in producing this report. The report has been through the SARDI internal review process, and has been formally approved for release by the Research Director, Aquatic Sciences. Although all reasonable efforts have been made to ensure quality, SARDI does not warrant that the information in this report is free from errors or omissions. SARDI and its employees do not warrant or make any representation regarding the use, or results of the use, of the information contained herein as regards to its correctness, accuracy, reliability and currency or otherwise. SARDI and its employees expressly disclaim all liability or responsibility to any person using the information or advice. Use of the information and data contained in this report is at the user's sole risk. If users rely on the information they are responsible for ensuring by independent verification its accuracy, currency or completeness. The SARDI Report Series is an Administrative Report Series which has not been reviewed outside the department and is not considered peer-reviewed literature. Material presented in these Administrative Reports may later be published in formal peer-reviewed scientific literature.

© 2021 SARDI

This work is copyright. Apart from any use as permitted under the *Copyright Act 1968* (Cth), no part may be reproduced by any process, electronic or otherwise, without the specific written permission of the copyright owner. Neither may information be stored electronically in any form whatsoever without such permission.

SARDI Publication No. F2020/000209-1
SARDI Research Report Series No. 1057

Author(s): T.M. Ward and G.L. Grammer

Reviewer(s): S. Mayfield (SARDI) and K. Bennetts (AFMA)

Approved by: S. Mayfield
Science Leader – Fisheries

Signed:



Date: 8 February 2021

Distribution: AFMA, SARDI Aquatic Sciences, Parliamentary Library, State Library and National Library

Circulation: Public Domain

Acknowledgements

This status summary report was funded by the Australian Fisheries Management Authority (AFMA). Data presented in this report were provided by: Ms Darci Wallis (Australian Fisheries Management Authority); Ms Denise Garcia and Ms Frances Seaborn (Department of Primary Industries, Parks, Water and Environment Tasmania); Dr John Stewart and Mr David Makin (New South Wales Department of Primary Industries); Ms Paula Baker and Ms Monique Nelis (Victorian Department of Economic Development, Jobs, Transport and Resources); and Ms Melleessa Boyle and Mr Angelo Tsolos (SARDI Aquatic Sciences). Catch samples were processed by Mr Alex Ivey, Mr Nat Navong and Mr Sam Foster. This report was formally reviewed and approved for release by Dr Stephen Mayfield (SARDI Aquatic Sciences).

Keywords: Commonwealth SPF, Jack Mackerel, Blue Mackerel, Redbait, Sardine, purse seine, mid-water trawl, AFMA.

Overview

This status summary report uses fishery-dependent and fishery-independent data to assess the status of target species in the East and West sub-areas of the Commonwealth Small Pelagic Fishery (SPF) in 2018/19. It updates the fishery assessment report of Ward and Grammer (2019).

The target species for the SPF are Jack Mackerel (*Trachurus declivis*), Blue Mackerel (*Scomber australasicus*), Redbait (*Emmelichthys nitidus*) and, in the Australian Sardine sub-area of the East sub-area, Australian Sardine (*Sardinops sagax*). The primary fishing methods are purse-seining and mid-water trawling. The assessment uses commercial catch and effort data for the SPF up to 30 April 2019 and biological information (size and age structures, reproduction, maturity). Data from State-managed commercial fisheries are used to estimate the total catch of each species in each sub-area.

The SPF Harvest Strategy 2008 (last revised April 2017) specifies that the primary technique for assessing the status of SPF species is the Daily Egg Production Method (DEPM). To retain a species in a sub-area at Tier 1, where exploitation rates are highest, the DEPM must be applied every 5 years. Between applications of the DEPM, fishery-dependent data are analysed to identify variations in fishing patterns or catches that may be indicative of changes in stock status.

Between 2010/11 and 2014/15, catches in the SPF of all species in both sub-areas were low (<200 t per annum). Effort and catch in the SPF increased during 2015/16 and 2016/17, when a mid-water factory trawler operated in offshore waters of both sub-areas. Since 2017/18, a smaller mid-water trawler without onboard processing facilities began operating in inshore waters of the East sub-area. Negligible fishing was undertaken by SPF vessels in the West sub-area in 2018/19.

Jack Mackerel

The estimate of spawning biomass for Jack Mackerel East in 2019 was 156,292 t (95% CI = 49,120–263,496). The total catch of Jack Mackerel East in 2018/19 was 4,947 t, the second highest catch in the last five years (highest: 6,321 t in 2015/16). The main fishery for Jack Mackerel East is the SPF. During 2018/19, the modal size of Jack Mackerel East taken in mid-water trawls (200 mm FL) was below the mean size at 50% maturity (~230 mm FL). Fluctuations in catches of Jack Mackerel East reflect changes in fishing effort. The total catch of Jack Mackerel East was 3% of the 2019 estimate of spawning biomass and below the Tier 1 exploitation rate for this stock of 12%. The total catch by the SPF in 2018/19 was 26% of the available TAC. On the basis of the information provided above, Jack Mackerel East is classified as **sustainable**.

The spawning biomass of Jack Mackerel West in 2016/17 was estimated to be at least 31,000 t. There was no catch of Jack Mackerel West by the SPF in 2018/19; this was

down from 686 t in 2016/17 and 634 t in 2015/16. The main fishery for Jack Mackerel West has been the SPF. Recent annual catches of Jack Mackerel West have been <3% of the 2016/17 estimate of spawning biomass and below the Tier 1 exploitation rate for this stock of 12%. The total catch in 2018/19 by the SPF was 0% of the available TAC. On the basis of the information provided above, Jack Mackerel West is classified as **sustainable**.

Blue Mackerel

The spawning biomass of Blue Mackerel East in 2014 was ~83,300 t (95% CI = 35,100–165,000 t). The total catch of Blue Mackerel East in 2018/19 was 4,265 t, which is the highest on record, eclipsing the previous maxima of 2,022 t in 2015/16 and 2,858 t in 2017/18. The main fisheries for Blue Mackerel East are the SPF (89% in 2018/19) and the New South Wales Ocean Hauling Fishery (11% in 2018/19). During 2018/19, the modal size of Blue Mackerel East from mid-water trawls (280 mm FL) was above the mean size at 50% maturity (~260 mm FL). Fluctuations in catches of Blue Mackerel East reflect changes in fishing effort. The total catch of Blue Mackerel East was 5% of the 2014 estimate of spawning biomass and below the Tier 1 exploitation rate for this stock of 15%. The total catch by the SPF in 2018/19 was 32% of the available TAC. On the basis of the information provided above, Blue Mackerel East is classified as **sustainable**.

The spawning biomass for Blue Mackerel West in 2005 was estimated to be 56,228 t. There was no catch of Blue Mackerel West by the SPF in 2018/19; this was down from 766 t in 2016/17 and 980 t in 2015/16. The main fishery for Blue Mackerel West has been the SPF. Recent total annual catches of Blue Mackerel West have been <2% of the estimated spawning biomass for 2005 and below the Tier 3 exploitation rate for this stock of 3.75%. The total catch in 2018/19 was 0% of the available TAC. On the basis of the information provided above, Blue Mackerel West is classified as **sustainable**.

Redbait

The spawning biomass of Redbait East was estimated to be ~70,000 t from surveys in 2005 and 2006. The total catch of Redbait East by the SPF in 2018/19 was 539 t, the highest annual catch in the last five years. The main fishery for Redbait East is the SPF. During 2018/19, the modal size of Redbait East from mid-water trawls (170 mm FL) was above the mean size at 50% maturity (~160 mm FL). Fluctuations in catches of Redbait East reflect changes in fishing effort. The total catch of Redbait East by the SPF was 1% of the estimate of spawning biomass and below the Tier 2 exploitation rate for this stock of 5%. The total catch by the SPF in 2018/19 was 16% of the available TAC. On the basis of the information provided above, Redbait East is classified as **sustainable**.

The estimate of spawning biomass for Redbait West in 2017 was 66,767 t (CI = 28,797–190,392). There was no catch of Redbait West by the SPF in 2018/19; this was down from 1,140 t in 2016/17 and 1,157 t in 2015/16. The main fishery for Redbait West has been the

SPF. Recent catches of Redbait West have been <2% of the 2017 estimate of spawning biomass and below the Tier 1 exploitation rate for this stock of 10%. The total catch in 2018/19 by the SPF was 0% of the available TAC. On the basis of the information provided above, Redbait West is classified as **sustainable**.

Australian Sardine

The estimate of spawning biomass of Australian Sardine in the Sardine sub-area in 2014 was 49,575 t (95% CI = 24,200–213,300 t). The total catch of Sardine in the Sardine sub-area in 2018/19 was 596 t; annual catches have been stable since 2013/14 (mean: 533 t·yr⁻¹; range: 429–601 t). The main fisheries for Sardine in the Sardine sub-area are the SPF (22% in 2018/19) and the New South Wales Ocean Hauling Fishery (78% in 2018/19). Catches of Sardine from the Sardine sub-area have mainly contained fish at or above the mean size at 50% maturity of ~150 mm FL. The total catch of Sardine in the Sardine sub-area was 1% of the 2014 estimate of spawning biomass and below the Tier 1 exploitation rate for this stock of 20%. The total catch by the SPF in 2018/19 was 1% of the available TAC. On the basis of the information provided above, Sardine in the Sardine sub-area is classified as **sustainable**.

Summary

All SPF stocks are classified as sustainable. DEPM surveys have now been conducted for all stocks. Resulting estimates of spawning biomass could be used to inform the establishment of target (e.g. B₅₀) and limit reference points (e.g. B₂₀) for each stock.

All SPF stocks are at Tier 1 except Blue Mackerel West and Redbait East. The DEPM was last applied to Blue Mackerel West in 2005. A DEPM survey of Blue Mackerel East and Australian Sardine in the Sardine sub-area was conducted in September 2019. A survey of Redbait East is scheduled for October 2020.

Reference

Ward, T. M. and Grammer, G. L. (2019). Commonwealth Small Pelagic Fishery: Fishery Assessment Report 2018. Report to the Australian Fisheries Management Authority. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2010/000270-10. SARDI Research Report Series No. 1021. 132pp.